



# Beast Exporter & Importer Pvt. Ltd.

*Export & Import Company*

Contact no. +91-8294983560, +91-7061189438

Add:- Bhagwati Nagar (Near; Harivatica chowck) bettiah, Bihar, pin 845438

Email id :- [beastexpoimpo.pvt.ltd@gmail.com](mailto:beastexpoimpo.pvt.ltd@gmail.com)

S.No	PRODUCT NAME	COMPOSITION	PACKING	MRP/BOX	NET RATE/BOX	PACK	HSN CODE	TAX%
<b>CARDIAC RANGE</b>								
1	ATOLVA-10	<b>Atorvastatin 10 mg</b> 1. Reduces the risk of myocardial infarction. 2. Atorvastatin is an oral drug that lowers the level of cholesterol in the blood. 3. Atorvastatin also reduces the concentration of triglycerides in the blood.	ALU-ALU	500.00	90.00	10x10	3004	12%
2	ATOLVA-20	<b>Atorvastatin 20 mg</b> 1. Reduces the risk of myocardial infarction. 2. Atorvastatin is an oral drug that lowers the level of cholesterol in the blood. 3. Atorvastatin als	ALU-ALU	1200.00	105.00	10x10	3004	12%
3	ATOLVA-C10	<b>Atorvastatin 10 mg + Clopidogrel 75 mg</b> 1. Effectively lowers plasma cholesterol & lipoprotein levels. 2. To prevent heart attacks & strokes in persons with heart disease (recent heart attack), recent stroke, or blood circulation disease (peripheral vascula	ALU-ALU	950.00	186.00	10x10	3004	12%
4	ATOLVA-F	<b>Atorvastatin 10mg + Fenofibrate 160mg</b> 1. Atorvastatin is used along with a proper diet to help lower "bad" cholesterol and fats (such as LDL, triglycerides) and raise "good" cholesterol (HDL) in the blood. 2. Fenofibrate helps reduc	10x10 TAB.	1800.00	195.00	10*10	3004	12%
5	AMNOP-AT	<b>Amlodipine 5mg + Atenolol 50mg</b> 1. A selective beta one blocker decreases plasma renin activity. 2. Inhibits the transmembrane influx of calcium ions vascular smooth muscle & cardiac muscle. 3. effective control of blood pressure also did not effct plasm	ALU-ALU	N.A	N.A	10x10	3004	12%
6	AMTOPROL-AL	<b>Metoprolol 50mg XL + Amlodipine 5mg</b> 1. well documented efficacy in angina pectoris reducing the frequency, duration & severity of angina attacks. 2. A selective calcium channel blocker vaso dilator. 3. Reduces total peripheral resistance.	BLISTER	750.00	180.00	10x10	3004	12%



17	<b>GLISPO-1M</b>	<b>Glimepride 1mg + Metformin 500mg (SR)</b> 1. Highly effective in patients having both high fasting and high post-prandial blood glucose levels. 2. The addition of glimepiride to MET is an effective treatment for patients inadequately controlled on sulfonylure	BLISTER (BI-LAYERED)	480.00	114.00	10x10	3004	12%
18	<b>GLISPO-2M</b>	<b>Glimepride 2mg + Metformin 500mg (SR)</b> After the intensive teatment, FPG,2 HPG & HBA1c decreased significantly(p>0.01); HOMA-IR desceased & HOMA-B incresed significantly (p<0.01) & TG,CHOL,LDL descreased significantly (p<0.01) after treatment	BLISTER (BI-LAYERED)	510.00	120.00	10x10	3004	12%
19	<b>GLISPO-1M FORTE</b>	<b>Glimepride 1mg + Metformin 1000mg (SR)</b> After the intensive teatment, FPG,2 HPG & HBA1c decreased significantly(p>0.01); HOMA-IR desceased & HOMA-B incresed significantly (p<0.01) & TG,CHOL,LDL descreased significantly (p<0.01) after treatment	BLISTER (BI-LAYERED)	N.A.	N.A.	10x10	3004	12%
20	<b>GLISPO-2M FORTE</b>	<b>Glimepride 2mg + Metformin 1000mg (SR)</b> After the intensive teatment, FPG,2 HPG & HBA1c decreased significantly(p>0.01); HOMA-IR desceased & HOMA-B incresed significantly (p<0.01) & TG,CHOL,LDL descreased significantly (p<0.01) after treatment	BLISTER (BI-LAYERED)	1000.00	168.00	10x10	3004	12%
21	<b>GLISPO-PM1/15</b>	<b>Glimepride 1mg + Metformin 500mg (SR) + Pioglitazone 15 mg</b> 1. Stimulates the release of insulin from functioning pancreatic beta cells. 2. Counters insulin resistance. 3. decreases heparic glucose production alo decreases intestinal absorption of glucose.	BLISTER (BI-LAYERED)	640.00	150.00	10x10	3004	12%

22	GLISPO-PM2/15	<p><b>Glimepride 2mg + Metformin 500mg (SR) + Pioglitazone 15 mg</b></p> <p>1. Stimulates the release of insulin from functioning pancreatic beta cells.  2. Counters insulin resistance.  3. decreases hepatic glucose production also decreases intestinal absorption of glucose</p>	BLISTER (BI-LAYERED)	910.00	158.00	10x10	3004	12%
23	TRIBOG-1.2	<p><b>Glimepride 1mg + Voglibose 0.2mg + Metformin 500mg (SR)</b></p> <p>1. Voglibose scores over both acarbose and miglitol in terms of side effect profile.  2. proven efficacy.  3. metformin causes few adverse effects the most common is gastrointestinal upset and is associated</p>	BLISTER (BI-LAYERED)	940.00	200.00	10x10	3004	12%
24	TRIBOG-2.2	<p><b>Glimepride 2mg + Voglibose 0.2mg + Metformin 500mg (SR)</b></p> <p>1. Voglibose scores over both acarbose and miglitol in terms of side effect profile. 2. proven efficacy.  3. metformin causes few adverse effects the most common</p>	BLISTER (BI-LAYERED)	1200.00	174.00	10x10	3004	12%
25	GLYCLIP-20	<p><b>Teneligliptin 20 mg</b></p> <p>Teneligliptin is used in the treatment of type 2 diabetes. It is used in addition to diet and exercise to improve blood sugar control in adults with type 2 diabetes.</p>	ALU-ALU	1250.00	216.00	10x10	3004	
26	GLYCLIP-M	<p><b>Teneligliptin 20 mg + Metformin 500 mg (PR)</b></p> <p>It is used in addition to diet and exercise to improve blood sugar control in adults with type 2 diabetes. Metformin is used with a proper diet and exercise program and possibly with other medications to control</p>	BLISTER (BI-LAYERED)	N.A.	N.A.	10x10	3004	12%

27	<b>LORIPTA-V</b>	<b>Vidagliptin 50mg</b> It helps to control the levels of sugar (glucose) in your blood. If your body does not make enough insulin, or if it does not use the insulin it makes effectively, this results in the condition called sugar diabetes	ALU-ALU	950.00	246.00	10*10	3004	12%
----	------------------	--	---------	--------	--------	-------	------	-----

28	<b>LORIPTA-V FORTE</b>	<b>Vidagliptin 100mg</b> It helps to control the levels of sugar (glucose) in your blood. If your body does not make enough insulin, or if it does not use the insulin it makes effectively, this results in the condition called sugar diabetes	ALU-ALU	1400.00	336.00	10*10	3004	12%
29	<b>LORIPTA-VM</b>	<b>Vidagliptin 50mg + Metformin 500 mg</b> It is used to treat type 2 diabetes mellitus in people who are already taking vildagliptin and metformin tablets separately, or whose diabetes cannot be controlled by metformin alone. ... Type 2 diabetes mellitus used t	BLISTER (BI-LAYERED)	1400.00	305.00	10*10	3004	12%
30	<b>LORIPTA-VM FORTE</b>	<b>Vidagliptin 50mg + Metformin 1000 mg</b> It is used to treat type 2 diabetes mellitus in people who are already taking vildagliptin and metformin tablets separately, or whose diabetes cannot be controlled by metformin alone. ... Type 2 diabetes mellitus used	BLISTER (BI-LAYERED)	1680.00	350.00	10*10	3004	12%
31	<b>VOBTRI-0.2MD</b>	<b>Voglibose 0.2mg (MD)</b> Voglibose scores over both acarbose and miglitol in terms of side effect profile.	ALU-ALU	N.A.	N.A.	10x10	3004	12%
32	<b>VOBTRI-0.3MD</b>	<b>Voglibose 0.3mg (MD)</b> Voglibose inhibitors delay glucose absorption at the intestine level and thereby prevent sudden surge of glucose after a meal.	ALU-ALU	880.00	135.00	10x10	3004	12%
33	<b>VOBTRI-0.2 MF</b>	<b>Voglibose 0.2 mg + Metformin 500 mg (SR)</b> Voglibose inhibitors delay glucose absorption at the intestine level and thereby prevent sudden surge of glucose after a meal.	BLISTER (BI-LAYERED)	850.00	189.00	10x10	3004	12%
34	<b>VOBTRI-0.3 MF</b>	<b>Voglibose 0.3 mg + Metformin 500 mg (SR)</b> Voglibose inhibitors delay glucose absorption at the intestine level and thereby prevent sudden surge of glucose after a meal.	BLISTER (BI-LAYERED)	N.A.	N.A.	10x10	3004	12%